

# Habitat-based density model for Kogia spp. in the AFTT area

*Laura Mannocci, Jason J Roberts, David L Miller, Patrick N Halpin*

*October 1, 2016*

---

This report documents the habitat-based density model for Kogia spp. in the Atlantic Fleet Testing and Training Area (AFTT) area. Information on the first stage of the modeling approach, including classification of ambiguous sightings, detection function fitting and  $g(0)$  estimation can be found in individual taxon reports presented in Roberts et al. (2016) for the U.S. Atlantic and Gulf of Mexico.

Citation for this model: Mannocci L, Roberts JJ, Miller DL, Halpin PN (2016). Habitat-based density model for Kogia spp. in the AFTT area. 2016-10-01. Marine Geospatial Ecology Lab, Duke University, Durham, NC.

Citation for the related publication: Mannocci L, Roberts JJ, Miller DL, Halpin PN. Extrapolating cetacean densities to quantitatively assess human impacts on populations in the high seas. In review in Conservation Biology.

## 1- Available data

Table 1: Effort (km) and sightings per surveyed region (CAR: Caribbean, EC: East coast, EU: European Atlantic, GM: Gulf of Mexico, MAR: Mid-Atlantic ridge). Details on the origin of sightings used in this study can be found in Table 1 of the associated publication.

Region	Effort	Sightings
CAR	24264.47	7
EC	1044357.70	31
GOM	194715.35	218
All regions	1263337.53	256

Table 2: Effort (km) and sightings per month.

Month	Effort	Sightings
January	77892.79	4
February	123591.37	4
March	117923.54	13
April	117929.72	14
May	149765.03	105
June	129393.69	29
July	135693.85	55
August	129660.43	24
September	71696.07	4
October	82560.18	1
November	69210.92	3
December	58019.93	0
All Months	1263337.53	256




Figure 1: Map of segments (black lines) and sighting locations (red dots). An Albers equal area projection optimized for the AFTT area is used.

## 2- Methodological decisions

Methodological decisions reported in this section were made according to information available to us in the literature as well as feedback from a number of experts we consulted.

### *Modeled taxon*

*Kogia* spp.

Since dwarf sperm whale (*Kogia sima*) and pygmy sperm whale (*Kogia breviceps*) are not differentiable at sea, leading to a high number of ambiguous sightings, we modeled them together within the *Kogia* spp. guild.

### *Modeled season*

We fitted a year-round model as we found no definitive evidence in the literature that *Kogia* undertakes extensive migrations or exhibits contrasting behaviors (e.g., feeding versus breeding) in different seasons at the scale of our study area.

### *Segments*

We incorporated segments from the east coast, Gulf of Mexico and Caribbean where sightings were reported (Table 1).

### 3- Best model

- **Predictors:** depth, production of epipelagic micronekton (EpiMnkPP), sea surface temperature (SST)
- **Model summary:**

```
##
## Family: Tweedie(p=1.312)
## Link function: log
##
## Formula:
## abundance ~ s(Depth, k = 4, bs = "ts") + s(EpiMnkPP, k = 4, bs = "ts") +
##           s(SST, k = 4, bs = "ts") + offset(log(area_km2))
## <environment: 0x1a61d35c>
##
## Parametric coefficients:
##               Estimate Std. Error t value Pr(>|t|)
## (Intercept)  -13.351      1.097  -12.17   <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Approximate significance of smooth terms:
##               edf Ref.df      F  p-value
## s(Depth)      2.248      3 19.665 4.43e-14 ***
## s(EpiMnkPP)   1.062      3  4.232 0.000215 ***
## s(SST)        1.697      3 21.753  < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## R-sq.(adj) =  -0.00228  Deviance explained = 41.8%
## -REML = 1986.3  Scale est. = 59.839    n = 113880
```




Figure 2: GAM term plots with the log-transformed abundance on the y axis. The solid blue line is the smooth function fitted to the data. The solid red line is the smooth function extrapolated to all covariate values in the prediction area. The dashed lines represent the approximate 95% confidence intervals. The rug plot on the x-axis shows covariate values sampled in the data. Note that transformations were used for some covariates.

## 4- Environmental envelopes






Figure 3: Environmental envelope for depth. White cells within the AFTT polygon indicate areas where covariate values fell beyond the range of covariate values sampled by the surveys.


**January**




**February**




**March**




**April**



**May**



**June**










Figure 4: Monthly environmental envelopes for production of epipelagic micronekton. White cells within the AFTT polygon indicate areas where covariate values fell beyond the range of covariate values sampled by the surveys.


**January**




**February**




**March**




**April**



**May**



**June**






Figure 5: Monthly environmental envelopes for sea surface temperature. White cells within the AFTT polygon indicate areas where covariate values fell beyond the range of covariate values sampled by the surveys.

## 5- Predicted densities




Figure 6: Mean predicted densities (individuals 100 km<sup>-2</sup>) in the AFTT area. Areas where we extrapolated beyond sampled predictor ranges and predicted densities should not be trusted are indicated with black crosshatches. An Albers equal area projection is used.

## 6- Coefficients of variation




Figure 7: Mean predicted coefficients of variation derived from GAM parameters in the AFTT area. Areas where we extrapolated beyond sampled predictor ranges and coefficients of variation should not be trusted are indicated with black crosshatches. An Albers equal area projection is used.

## 7- Predicted densities per province




Figure 8: Predicted densities (individuals 100 km<sup>2</sup>) averaged per Longhurst's biogeographical province. Note that the color scheme is the same as in Figure 7. Provinces: ARCT: Atlantic Arctic Province; BPLR: Boreal Polar Province; CARB: Caribbean Province; GFST: Gulf Stream Province; NATR: North Atlantic Tropical Gyral Province; NASW: North Atlantic Subtropical Gyral Province (West); NWCS: North West Atlantic Shelves Province.

## 8- Alternate models




Figure 10: Sensitivity of densities predicted by the five top models per Longhurst's biogeographical province. Points represent predicted densities (individuals 100 km<sup>2</sup>) for the five top models listed in Table 3, with the first to fifth models ordered from left to right. Filled points correspond to models with some support (*sensu* Burnham and Anderson (2002), i.e.,  $\Delta AIC < 2$ ) while hollow points correspond to models with little support (i.e.,  $\Delta AIC > 2$ ). The shaded areas indicate the range of densities predicted by the five top models for each province. Provinces: ARCT: Atlantic Arctic Province; BPLR: Boreal Polar Province; CARB: Caribbean Province; GFST: Gulf Stream Province; NATR: North Atlantic Tropical Gyral Province; NASW: North Atlantic Subtropical Gyral Province (West); NWCS: North West Atlantic Shelves Province.

Table 3: List of the five top models with lowest AIC values. Ns: non-significant. Predictor variables: EKE: eddy kinetic energy, SLAStDev: standard error of sea level anomaly, SST: sea surface temperature, PkPP: zooplankton production, PkPB: zooplankton biomass, EpiMnkPP: epipelagic micronekton production, EpiMnkPB: epipelagic micronekton biomass, VGPM: vertically generalized production model, CHL: chlorophyll-a concentration.

		Predictors		AIC	delta AIC
Depth	SST	EpiMnkPP	ns	113959.3	0.0
Depth	SST	EpiMnkPB	ns	113960.8	1.5
Depth	DistToFront1	EpiMnkPB	ns	113961.0	1.7
Depth	SST	EpiMnkPB	SLAStDev	113961.7	2.4
Depth	SST	EpiMnkPP	SLAStDev	113962.5	3.2



## 9- Residual diagnostics




Figure 10: Diagnostic plots of residuals. Left: Quantile-quantile (Q-Q) plot of deviance residuals generated using the `qq.gam` function with 100 simulations (Augustin et al. 2012). Grey lines are possible simulated Q-Q plots under the assumption that the model is correct. The red reference line indicates perfect agreement between residual and theoretical residual distributions. Points lying away from the red line suggest poor model fit for the corresponding quantiles. Zeros appear to the left of the Q-Q plot in alignment with the reference line. Because, by design, models were not tightly fitted to the data (see discussion of the paper), deviations from the red line may be observed. Specifically, points far above the red line for large quantiles indicate that the model underestimates high abundances observed on some segments. Right: randomized quantile residuals vs. linear predictor. A LOWESS regression is shown as a red line to illustrate any trend in the points. This plot should be generally free of any pattern. Expanding y-range indicates non-constant variance (heteroskedasticity) in the model.




Figure 11: Boxplots of deviance residuals, binned for each predictor. The horizontal line represents the median, and the bottom and top of the box represent the first and third quartiles respectively. Whiskers extend 1.5 times the inter-quartile range following McGill et al. (1978). Total counts of outliers beyond the whiskers are indicated in red. Numbers of segments per bin are indicated in grey. Boxplots for the different bins of predictors should generally overlap. A boxplot having its median away from zero indicates poorer model fit for that predictor bin. Boxplots often have their medians close to zero and fewer outliers for predictor bins characterized by low abundances of the species, suggesting that model fit is generally better in low abundance areas. We believe this is an inherent feature of models applied to count data with numerous zeros.

## 10- Brief discussion and overall confidence in predictions

### *Description of confidence levels*

We group taxa in three categories reflecting our relative level of confidence in predicted densities.

#### Level 1

This category includes tropical and warm temperate taxa for which survey data were available within most of the distributional range in the AFTT area. High/intermediate densities predicted beyond surveyed areas were supported by sightings available from OBIS-SEAMAP and the scientific literature. Very low densities predicted at northern latitudes were consistent with the described absence of these taxa. We have a reasonable confidence in predicted densities for these taxa.

#### Level 2

This category encompasses taxa for which a large part of the distributional range is in cold temperate and sub-polar waters. Models fitted to available survey data and extrapolated to cold temperate and sub-polar waters successfully predicted their occurrence, but predicted densities were largely speculative. The incorporation of line transect survey data from Canada and Greenland would be extremely useful to increase the reliability of predicted densities at northern latitudes. Unfortunately we were unable to obtain permission for using these data in our models. We remain hopeful that collaborations can be established in the future, and that the Canadian and Greenlandic surveys may be incorporated into a new version of our models. We have medium or low confidence in predicted densities for these taxa.

#### Level 3

This category includes taxa that are not known to primarily occur in cold temperate and sub-polar waters but were predicted in low/intermediate densities at higher latitudes. For these taxa, we believe predicted densities were likely overestimated at higher latitudes. However, predicted densities were supported by sightings available from OBIS-SEAMAP and the scientific literature within their core distributional range. The incorporation of line transect survey data from Canada and Greenland would be extremely useful to help correct the probable overestimation of densities at northern latitudes. We remain hopeful that collaborations can be established in the future, and that the Canadian and Greenlandic surveys may be incorporated into a new version of our models. We have medium or low confidence in predicted densities for these taxa.




Figure 12: Red points are sightings of the taxon from line transect surveys used in this study. Blue points are sightings of the taxon reported by other datasets not used in our study for 1992-2016 (e.g., because they were not compatible with our methodology). Underlain grey points are sightings of other cetacean species, taken from these other datasets. Blue and grey points were extracted from OBIS-SEAMAP (accessible at <http://seamap.env.duke.edu/>) (Halpin et al. 2009); citations for individual datasets are provided at the end of this report. Longhurst's biogeographical provinces are shown as polygons. Dense patches of grey points without red or blue points suggest locations where the taxon of interest may be absent, under the presumption that observers who reported other cetacean taxa would have reported this one if sighted. However, important caveats apply: the map does not quantify observation effort, which was not available for all datasets and was very difficult to standardize across disparate sources (e.g., scientific surveys, whale watching logs, opportunistic sightings). The spatial distribution of effort was highly heterogeneous in both space and time. Only openly accessible datasets were considered; other cetacean datasets are known to exist for the AFTT area but have not been released for public use (e.g., the 2007 Trans North Atlantic Sightings Survey (TNASS) in Canada). The presumption that grey dots imply absence may not always hold; for example, if effort conducted in that area was directed towards particular species, sightings of our taxon of

interest may not have been recorded.

### *General*

A total of 256 sightings, mostly from surveys in the northern Gulf of Mexico, were available to fit the habitat-based density model. The first or lowest AIC model included sea surface temperature, depth and epipelagic micronekton production (listed in decreasing order of importance according to F-scores) and had an explained deviance of 41.8%. All top five models included depth (Table 3). All but the third model included sea surface temperature. The second and third models had a delta AIC  $< 2$ , indicating some statistical support sensu Burnham and Anderson (2002). The fourth and fifth models had comparatively little support. Predicted densities from the top five models were quite variable in the different provinces (Figure 9). The third model was largely responsible for this variation, predicting consistently higher densities than the other models. In the ARCT and BPLR provinces, well beyond the northern distribution range of *Kogia* spp. (Bloodworth & Odell 2008; McAlpine 2009), all models predicted densities near zero except the third model. Despite having some statistical support, the third model led to unrealistically high predicted densities so we decided to exclude it. Predicted densities from the first and second statistically supported models were overall similar in all provinces. The first model had a lower AIC and explained more deviance compared to the second model, suggesting it was slightly more suitable for modeling *Kogia* spp. densities.

Model predictions appeared generally consistent with the described range of *Kogia* spp. in offshore tropical and temperate waters (there is some evidence that dwarf sperm whale prefers warmer waters than pygmy sperm whale) (Bloodworth & Odell 2008; McAlpine 2009). The higher predicted densities along the continental slope appeared in line with habitat preferences of *Kogia* spp. inferred from diet analyses of stranded animals (McAlpine et al. 2009). The scarcity of sightings reported beyond well surveyed waters is likely due to the cryptic behavior and difficulty in identifying these species at sea.

We now discuss the quality of predictions per biogeographic province by comparing them with the available literature and observations from OBIS-SEAMAP.

### *Boreal polar (BPLR) and Atlantic Arctic (ARCT) provinces*

Low predicted densities in the BPLR province, the ARCT province and the northern part of the NWCS province (approx. north of 42°N) seemed consistent with the described distribution of *Kogia* spp. in tropical and temperate waters (Bloodworth & Odell 2008; McAlpine 2009). We note that a few extralimital records exist in eastern Canada (McAlpine et al. 1997) and the northernmost sighting reported in OBIS-SEAMAP was 43°N. We caution that extrapolation to colder waters occurred in the BPLR, ARCT and northern part of the NWCS province, potentially leading to unreliable predictions. Nevertheless, we believe the low predicted densities in these northern waters are in line with the ecology of *Kogia* spp.

### *North West Atlantic shelves (NWCS) and Gulf Stream (GFST) provinces*

Predicted densities were the highest in offshore waters south of Cape Hatteras, North Carolina, where *Kogia* were recorded by multiple surveys. North of Cape Hatteras, *Kogia* were also recorded by surveys and numerous sightings were available in OBIS-SEAMAP (Figure 12), but predicted densities were lower. We think there is a possibility that our model underestimates *Kogia* densities north of Cape Hatteras.

### *North Atlantic tropical gyral (NATR) and North Atlantic subtropical gyral (NASW) provinces*

The model predicted medium densities in the NATR and NASW provinces. Only one offshore sighting was reported in the NASW province and no sighting was reported in the NATR province, but observation effort in these provinces was extremely sparse (Figure 12). Predictions near the Bahamas (edge of the NASW province) appeared consistent with sightings of dwarf sperm whales east of Great Abaco where they occur at relatively low depths community (MacLeod et al. 2004) (Figure 12).

Until they are validated with additional sightings, predictions in offshore waters of the NATR and NASW provinces will remain largely speculative.

### *Caribbean (CARB) province*

The CARB province was the province where predicted density was the highest on average.

In the Gulf of Mexico, predicted densities were the highest on the continental slope. Pygmy and dwarf sperm whales are considered common species within the offshore cetacean fauna of the Gulf of Mexico (Ortega-Ortiz 2002). Little information exists for *Kogia* spp. in the southern Gulf of Mexico. Ortega-Ortiz (2002) reported 2 sightings from an opportunistic cruise and 15 strandings on the Mexican coast. Given the numerous sightings in offshore waters of the northern Gulf of Mexico, we believe predicted densities in offshore waters of the southern Gulf of Mexico are not unrealistic.

We note that 13 strandings of *Kogia* spp. were reported for Puerto Rico and the Virgin Islands (Cardona-Maldonado and Mignucci-Giannoni 1999).

*Overall confidence: level 1*

Large amounts of survey data were available within the core distributional range of *Kogia* spp. and predicted densities seemed compatible with their described pelagic ecology. Few sightings were available to support predicted densities beyond surveyed areas but we believe this paucity of sightings is largely due to the cryptic behavior characterizing this taxon. The model predicted very low densities in northern waters, consistent with the described absence of *Kogia* spp.

## 11- References

- Augustin NH, Sauleau E-A, Wood SN. 2012. On quantile quantile plots for generalized linear models. *Computational Statistics & Data Analysis* 56:2404-2409.
- Burnham KP, Anderson DR. 2002. *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*. Springer Science & Business Media.
- Bloodworth BE, Odell DK. 2008. *Kogia breviceps* (Cetacea: Kogiidae). *Mammalian Species* 819:1-12.
- Cardona-Maldonado, M. A., and A. A. Mignucci-Giannoni. 1999. Pygmy and dwarf sperm whales in Puerto Rico and the Virgin Islands, with a review of *Kogia* in the Caribbean. *Caribbean Journal of Science* 35:29-37.
- Halpin P et al. 2009. OBIS-SEAMAP: The World Data Center for Marine Mammal, Sea Bird, and Sea Turtle Distributions. *Oceanography* 22:104-115.
- McAlpine DF. 2009. Pygmy and dwarf sperm whales. Pages 936-938 *Encyclopedia of marine mammals* 2nd Edition. Academic Press.
- McAlpine DF, Murison LD, Hoberg EP. 1997. New Records for the Pygmy Sperm Whale, *Kogia Breviceps* (physeteridae) from Atlantic Canada with Notes on Diet and Parasites. *Marine Mammal Science* 13:701-704.
- MacLeod CD, Hauser N, Peckham H. 2004. Diversity, relative density and structure of the cetacean community in summer months east of Great Abaco, Bahamas. *Journal of the Marine Biological Association of the United Kingdom* 84:469-474.
- McGill R, Tukey JW, Larsen WA. 1978. Variations of Box Plots. *The American Statistician* 32:12-16.
- Ortega-Ortiz, J. 2002. Multiscale analysis of cetacean distribution in the Gulf of Mexico. PhD dissertation Texas A&M University. Roberts JJ et al. 2016. Habitat-based cetacean density models for the U.S. Atlantic and Gulf of Mexico. *Scientific Reports* 6:22615.

### Citations for individual datasets from OBIS-SEAMAP

- Ampela, K. and G. Miller-Francisco. 2016. JAX FIREX Aerial Surveys 5-8 September 2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/880>) on 2016-08-15.
- Barco, S. 2014. Virginia and Maryland Sea Turtle Research and Conservation Initiative Aerial Survey Sightings, May 2011 through July 2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1201>) on 2016-08-15.
- Barco, S. 2014. Virginia CZM Wind Energy Area Survey- Vessel Survey Sightings - November 2012 through April 2014. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1196>) on 2016-08-15.
- Barco, S. 2015. Marine Mammal and Sea Turtle Sightings in the Vicinity of the Maryland Wind Energy Area 2013-2015. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1340>) on 2016-08-15.
- Barco, S. 2015. Virginia CZM Wind Energy Area Survey - Left side - May 2014 through December 2014. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1229>) on 2016-08-15.
- Barco, S. 2015. Virginia CZM Wind Energy Area Survey - Right side - May 2014 through December 2014. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1231>) on 2016-08-15.
- Barco, S. 2015. Virginia CZM Wind Energy Area Survey- Right side - November 2012 through April 2014. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1194>) on 2016-08-15.
- Barco, S. 2016. Virginia CZM Wind Energy Area Survey- Left side - November 2012 through April 2014. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1192>) on 2016-08-15.
- Boisseau, O. 2014. Visual sightings from Song of the Whale 1993-2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1158>) on 2016-08-15.
- Bolanos, J., J. Blumenthal, J. Luksenburg, A. Henriquez, A. Bogomolni, A. Mignucci-Giannoni, N. Landrau, J. Casas, M. Iniguez, J. Khan, C. Rinaldi, G. Ferrer, L. Suttty and N. Ward. 2014. Killer whales of the Caribbean

Sea 1866-2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1012>) on 2016-08-15.

Cole, T. and C. Khan. 2016. NEFSC Right Whale Aerial Survey. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/513>) on 2016-08-15.

Contillo, J. 2013. SEFSC Dolphin Photo ID. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/226>) on 2016-08-15.

Diaz, G. 2011. NOAA Southeast Fishery Science Center (SEFSC) Commercial Pelagic Observer Program (POP) Data. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/103151496>) on 2016-08-15 and originated from iOBIS (<http://www.iobis.org>).

DiMatteo, A. 2013. US Navy marine mammal and sea turtle sightings from aerial surveys, Vieques, Puerto Rico 2000. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1007>) on 2016-08-15.

Dunn, C. 2013. Bahamas Marine Mammal Research Organisation Opportunistic Sightings. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/329>) on 2016-08-15.

Epperson, D. 2012. BOEM Sperm Whale Seismic Study (SWSS) S-Tag Argos Telemetry. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/810>) on 2016-08-15.

Epperson, D. 2013. BOEM Sperm Whale Seismic Study (SWSS) MPS cetacean sightings 2002-2004. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/911>) on 2016-08-15.

Epperson, D. 2013. BOEM Sperm Whale Seismic Study (SWSS) MPS sperm whale trackings 2004-2005. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/912>) on 2016-08-15.

Epperson, D. 2013. BOEM Sperm Whale Seismic Study (SWSS) PhotoID 2002-2005. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/910>) on 2016-08-15.

Epperson, D. 2013. BOEM Sperm Whale Seismic Study (SWSS) S-Tag cetacean sightings 2002-2004. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/903>) on 2016-08-15.

Epperson, D. 2013. BOEM Sperm Whale Seismic Study (SWSS) S-Tag cetacean sightings 2005. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/905>) on 2016-08-15.

Epperson, D. 2015. BOEM Sperm Whale Seismic Study (SWSS) S-Tag sperm whale trackings 2002-2004. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/904>) on 2016-08-15.

Garrison, L. 2013. Gomex Sperm Whale Survey 2000. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/96>) on 2016-08-15.

Garrison, L. 2013. SEFSC Atlantic surveys 1992. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/3>) on 2016-08-15.

Garrison, L. 2013. SEFSC Atlantic surveys 1999. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/5>) on 2016-08-15.

Garrison, L. 2013. SEFSC Atlantic surveys, 1998 (3). Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1>) on 2016-08-15.

Garrison, L. 2013. SEFSC Caribbean Survey 1995. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/11>) on 2016-08-15.

Garrison, L. 2013. SEFSC Caribbean Survey 2000. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/7>) on 2016-08-15.

Garrison, L. 2013. SEFSC GoMex Oceanic 1992 (199). Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/13>) on 2016-08-15.

Garrison, L. 2013. SEFSC GoMex Oceanic 1993 (S). Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/17>) on 2016-08-15.

Garrison, L. 2013. SEFSC GoMex Oceanic 1993 (W). Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/15>) on 2016-08-15.

Garrison, L. 2013. SEFSC GoMex Oceanic 1994. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/19>) on 2016-08-15.

Garrison, L. 2013. SEFSC GoMex Oceanic 1996. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/25>) on 2016-08-15.

Garrison, L. 2013. SEFSC GoMex Oceanic 1997. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/27>) on 2016-08-15.

Garrison, L. 2013. SEFSC GoMex Oceanic 1999. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/29>) on 2016-08-15.

Garrison, L. 2013. SEFSC GoMex Oceanic 2000. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/29>) on 2016-08-15.



[env.duke.edu/dataset/21](http://env.duke.edu/dataset/21)) on 2016-08-15.

Garrison, L. 2013. SEFSC GoMex Oceanic 2001. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/23>) on 2016-08-15.

Garrison, L. 2013. SEFSC Gomex Shelf 1994. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/35>) on 2016-08-15.

Garrison, L. 2013. SEFSC Gomex Shelf 1998. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/31>) on 2016-08-15.

Garrison, L. 2013. SEFSC Gomex Shelf 2000. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/37>) on 2016-08-15.

Garrison, L. 2013. SEFSC Gomex Shelf 2001. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/33>) on 2016-08-15.

Garrison, L. 2013. SEFSC Mid-Atlantic Tursiops Survey, 1995 (1). Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/90>) on 2016-08-15.

Garrison, L. 2013. SEFSC Mid-Atlantic Tursiops Survey, 1995 2. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/89>) on 2016-08-15.

Garrison, L. 2013. SEFSC Mid-Atlantic Tursiops Survey, 1995 3. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/88>) on 2016-08-15.

Garrison, L. 2013. SEFSC Southeast Cetacean Aerial Survey 1992. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/87>) on 2016-08-15.

Garrison, L. 2013. SEFSC Southeast Cetacean Aerial Survey 1995. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/86>) on 2016-08-15.

Harris, Lei E. 2015. DFO Maritimes Region Cetacean Sightings. Version 6 In OBIS Canada Digital Collections. Bedford Institute of Oceanography, Dartmouth, NS, Canada. Published by OBIS, Digital <http://www.iobis.org/>.

Holst, M., O. Lee and H. Smith. 2014. Lamont-Doherty/LGL/NSF cruises. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/511>) on 2016-08-15.

Hyrenbach, D. 2011. Hatteras Eddy Cruise 2004. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/322>) on 2016-08-15.

Hyrenbach, D. and H. Whitehead. 2008. Sargasso sperm whales 2004. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/306>) on 2016-08-15.

Hyrenbach, D. and H. Whitehead. 2013. Sargasso 2005 - cetacean sightings. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/332>) on 2016-08-15.

Hyrenbach, D., F. Huettmann and J. Chardine. 2012. PIROP Northwest Atlantic 1965-1992. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/280>) on 2016-08-15.

Johnston, D. and Z. Swaim. 2013. DUML vessel-based surveys for proposed JAX USWTR site 2009-2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/582>) on 2016-08-15.

Josephson, B. 2015. AMAPPS Northeast Aerial Cruise Fall 2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1245>) on 2016-08-15.

Josephson, B. 2015. AMAPPS Northeast Aerial Cruise Spring 2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1247>) on 2016-08-15.

Josephson, B. 2015. AMAPPS Northeast Aerial Cruise Summer 2010. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1249>) on 2016-08-15.

Josephson, B. 2015. AMAPPS Northeast Aerial Cruise Summer 2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1233>) on 2016-08-15.

Josephson, B. 2015. AMAPPS Northeast Aerial Cruise Winter 2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1243>) on 2016-08-15.

Josephson, B. 2015. AMAPPS Northeast Shipboard Cruise Summer 2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1269>) on 2016-08-15.

Josephson, B. 2015. AMAPPS Northeast Shipboard Cruise Summer 2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1271>) on 2016-08-15.

Josephson, B. 2016. AMAPPS Northeast Aerial Cruise Spring 2014. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1379>) on 2016-08-15.

Josephson, B. 2016. AMAPPS Northeast Aerial Cruise Winter 2014. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1381>) on 2016-08-15.

Josephson, B. 2016. AMAPPS Northeast Shipboard Cruise Spring 2014. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1377>) on 2016-08-15.

Josephson, B. and L. Garrison. 2015. AMAPPS Southeast Aerial Cruise Fall 2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1288>) on 2016-08-15.

Josephson, B. and L. Garrison. 2015. AMAPPS Southeast Aerial Cruise Spring 2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1259>) on 2016-08-15.

Josephson, B. and L. Garrison. 2015. AMAPPS Southeast Aerial Cruise Summer 2010. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1273>) on 2016-08-15.

Josephson, B. and L. Garrison. 2015. AMAPPS Southeast Aerial Cruise Summer 2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1275>) on 2016-08-15.

Josephson, B. and L. Garrison. 2015. AMAPPS Southeast Aerial Cruise Winter 2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1277>) on 2016-08-15.

Josephson, B. and L. Garrison. 2015. AMAPPS Southeast Aerial Cruise Winter 2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1289>) on 2016-08-15.

Kopelman, A. 2013. Opportunistic marine mammal sightings from commercial whale watching vessels, Montauk, New York 1981-1994. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1006>) on 2016-08-15.

Kopelman, A. 2015. CRESLI marine mammal observations from whale watch cruises 2000-2014. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/896>) on 2016-08-15.

LaBrecque, E. 2011. Cape Hatteras 04-05. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/298>) on 2016-08-15.

Lanfredi, C. and G. Notarbartolo di Sciara. 2014. Tethys Research Institute shipboard survey cetacean sightings 1986-2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/774>) on 2016-08-15.

Lapolla, F. 2013. The Dolphin Project. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/304>) on 2016-08-15.

Latusek-Nabholz, J. 2013. Sightings for Airborne Mine Neutralization System Aerial Monitoring in the NSWC PCD Study Area from October 2011, 2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/948>) on 2016-08-15.

Latusek-Nabholz, J. 2013. Sightings for Airborne Mine Neutralization System Vessel Monitoring in the NSWC PCD Study Area from December 2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/949>) on 2016-08-15.

Latusek-Nabholz, J. 2013. Sightings for AN/AQS-20 Sonar Aerial Monitoring in the NSWC PCD Study Area from July 2011 and May 2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/947>) on 2016-08-15.

Latusek-Nabholz, J. 2014. Acoustic Detections for Airborne Mine Neutralization System Passive Acoustic Monitoring in the NSWC PCD Study Area from December 2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/950>) on 2016-08-15.

Latusek-Nabholz, J. 2014. Sightings for AN/AQS-20 Sonar Test Event - April-May 2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1096>) on 2016-08-15.

Latusek-Nabholz, J. 2014. Sightings for AN/AQS-20 Sonar Test Event - December 2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1113>) on 2016-08-15.

Latusek-Nabholz, J. 2014. Sightings for REMUS Sonar Test Event - July 2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1098>) on 2016-08-15.

Latusek-Nabholz, J. 2014. Sightings for SSAM2-BOSS Sonar Test Event - June 2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1097>) on 2016-08-15.

Mallette S.D., Lockhart G.G., McAlarney R.J., Cummings E.W., Pabst D. A., McLellan W.A., Barco S.G. 2016. Offshore Energy Planning: Documenting Megafauna off Virginia's Coast Using Aerial Surveys. VAQF Scientific Report. 2016-04.

Mallette S.D., Lockhart G.G., McAlarney R.J., Cummings E.W., Pabst D. A., McLellan W.A., Barco S.G. 2016. Offshore Energy Planning: Documenting Megafauna off Virginia's Coast Using Aerial Surveys. VAQF Scientific Report. 2016-04.

Maughan, B. and K. Arnold. 2010. UK Royal Navy Marine Mammal Observations. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/64>) on 2016-08-15.

McLellan, W. 2005. UNCW Aerial Survey 1998-1999. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/272>) on 2016-08-15.

McLellan, W. 2006. UNCW Marine Mammal Sightings 1998-1999. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/66>) on 2016-08-15.

McLellan, W. 2007. UNCW Marine Mammal Sightings 2002. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/67>) on 2016-08-15.

McLellan, W. 2010. UNCW Marine Mammal Sightings 2001. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/65>) on 2016-08-15.

McLellan, W. 2011. UNCW Aerial Surveys for monitoring of proposed Onslow Bay USWTR site - Left side -. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/435>) on 2016-08-15.

McLellan, W. 2011. UNCW Marine Mammal Aerial Surveys 2006-2007. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/400>) on 2016-08-15.

McLellan, W. 2011. UNCW Right Whale Aerial Survey 05-06. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/360>) on 2016-08-15.

McLellan, W. 2011. UNCW USWTR JAX Aerial Surveys May - Oct 2010 - Left side. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/687>) on 2016-08-15.

McLellan, W. 2011. UNCW USWTR JAX Aerial Surveys May - Oct 2010 - Right side. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/688>) on 2016-08-15.

McLellan, W. 2011. USWTR JAX Aerial Survey -Left side- 2009-2010. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/590>) on 2016-08-15.

McLellan, W. 2011. USWTR JAX Aerial Survey -Left side- 2010-2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/745>) on 2016-08-15.

McLellan, W. 2011. USWTR JAX Aerial Survey -Right side- 2010-2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/747>) on 2016-08-15.

McLellan, W. 2011. USWTR Onslow Bay Aerial Survey -Left side- 2008-2010. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/586>) on 2016-08-15.

McLellan, W. 2011. USWTR Onslow Bay Aerial Survey -Left side- 2010-2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/749>) on 2016-08-15.

McLellan, W. 2011. USWTR Onslow Bay Aerial Survey -Right side- 2008-2010. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/588>) on 2016-08-15.

McLellan, W. 2011. USWTR Onslow Bay Aerial Survey -Right side- 2010-2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/751>) on 2016-08-15.

McLellan, W. 2012. USWTR JAX Aerial Survey -Left side- 2011-2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/857>) on 2016-08-15.

McLellan, W. 2012. USWTR JAX Aerial Survey -Right side- 2009-2010. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/592>) on 2016-08-15.

McLellan, W. 2012. USWTR JAX Aerial Survey -Right side- 2011-2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/859>) on 2016-08-15.

McLellan, W. 2013. UNCW Aerial Surveys for monitoring of proposed Onslow Bay USWTR site - Right side -. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/437>) on 2016-08-15.

McLellan, W. 2013. UNCW Right Whale Aerial Surveys 2008. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/464>) on 2016-08-15.

McLellan, W. 2014. AFAST Hatteras Aerial Survey -Left side- 2011-2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/851>) on 2016-08-15.

McLellan, W. 2014. AFAST Hatteras Aerial Survey -Right side- 2011-2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/855>) on 2016-08-15.

McLellan, W. 2014. AFTT Hatteras Aerial Survey -Left side- 2012-2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1138>) on 2016-08-15.

McLellan, W. 2014. AFTT Hatteras Aerial Survey -Right side- 2012-2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1140>) on 2016-08-15.

McLellan, W. 2014. AFTT JAX Aerial Survey -Left side- 2012-2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1128>) on 2016-08-15.

McLellan, W. 2014. AFTT JAX Aerial Survey -Right side- 2012-2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1136>) on 2016-08-15.

McLellan, W. 2015. AFTT Cape Hatteras Aerial Survey -Left side- 2014. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1237>) on 2016-08-15.

McLellan, W. 2015. AFTT Cape Hatteras Aerial Survey -Right side- 2014. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1235>) on 2016-08-15.

McLellan, W. 2015. AFTT JAX Aerial Survey -Left side- 2014. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1241>) on 2016-08-15.

McLellan, W. 2015. AFTT JAX Aerial Survey -Right side- 2014. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1239>) on 2016-08-15.

McLellan, W. 2016. UNCW Hatteras Aerial Survey - Left side - 2015. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1350>) on 2016-08-15.

McLellan, W. 2016. UNCW Hatteras Aerial Survey - Right side - 2015. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1352>) on 2016-08-15.

McLellan, W. 2016. UNCW JAX Aerial Survey - Left side - 2015. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1362>) on 2016-08-15.

McLellan, W. 2016. UNCW JAX Aerial Survey - Right side - 2015. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1364>) on 2016-08-15.

McLellan, W. 2016. UNCW Norfolk Canyon Aerial Survey - Left side - 2015. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1354>) on 2016-08-15.

McLellan, W. 2016. UNCW Norfolk Canyon Aerial Survey - Right side - 2015. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1356>) on 2016-08-15.

Olsen, E. 2013. Long-Distance Movement of a Sei Whale in the North Atlantic, 2005. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/888>) on 2016-08-15.

Palka, D. 2011. NEFSC 1995 AJ9501 (Part I). Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/56>) on 2016-08-15.

Palka, D. 2013. Harbor Porpoise Survey 1992 (AJ92-01). Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/302>) on 2016-08-15.

Palka, D. 2013. NEFSC 1995 AJ9501 (Part II). Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/290>) on 2016-08-15.

Palka, D. 2013. NEFSC 1995 pe9501. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/296>) on 2016-08-15.

Palka, D. 2013. NEFSC 1995 pe9502. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/294>) on 2016-08-15.

Palka, D. 2013. NEFSC 1999 aj9902. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/300>) on 2016-08-15.

Palka, D. 2013. NEFSC Aerial Circle-Back Abundance Survey 2004. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/398>) on 2016-08-15.

Palka, D. 2013. NEFSC Aerial Survey - Experimental 2002. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/107>) on 2016-08-15.

Palka, D. 2013. NEFSC Aerial Survey - Summer 1995. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/109>) on 2016-08-15.

Palka, D. 2013. NEFSC Aerial Survey - Summer 1998. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/113>) on 2016-08-15.

Palka, D. 2013. NEFSC Deepwater Marine Mammal 2002. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/292>) on 2016-08-15.

Palka, D. 2013. NEFSC Mid-Atlantic Marine Mammal Abundance Survey 2004. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/396>) on 2016-08-15.

Palka, D. 2013. NEFSC Survey 1997. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/58>) on 2016-08-15.

Palka, D. 2013. NEFSC Survey 1998 1. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/60>) on 2016-08-15.

Palka, D. 2013. NEFSC Survey 1998 2. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/62>) on 2016-08-15.

Read, A. 2012. Duke Harbor Porpoise Tracking. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/83>) on 2016-08-15.



Serrano, A. 2011. Cetacean diversity, distribution, and abundance in northern Veracruz, Mexico. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/530>) on 2016-08-15.

Smith, A. 2014. Mystic Aquarium's marine mammal and sea turtle stranding data 1976-2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/945>) on 2016-08-15.

Speakman, T. 2011. NOAA Atlantic bottlenose dolphin sightings in the coastal and estuarine waters near Charleston, SC - 1994-2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/737>) on 2016-08-15.

Spontak, D. 2012. JAX ASWEX Aerial Monitoring 2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/868>) on 2016-08-15.

Spontak, D. 2012. JAX MAVEX Aerial Monitoring 2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/875>) on 2016-08-15.

Spontak, D. 2012. JAX MISSILEX Aerial Monitoring 2010. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/874>) on 2016-08-15.

Spontak, D. 2012. JAX SEASWITI Aerial Monitoring 2010 . Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/866>) on 2016-08-15.

Spontak, D. 2012. JAX SEASWITI Vessel Monitoring 2010. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/867>) on 2016-08-15.

Spontak, D. 2012. VACAPES ASWEX Aerial Monitoring 2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/869>) on 2016-08-15.

Spontak, D. 2012. VACAPES FIREX Aerial Monitoring 2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/871>) on 2016-08-15.

Spontak, D. 2013. JAX GUNEX Aerial Monitoring Surveys October 2010. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/893>) on 2016-08-15.

Spontak, D. 2013. JAX MAVEX September 2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/895>) on 2016-08-15.

Spontak, D. 2013. VACAPES FIREX and ASW Aerial Monitoring 2010. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/870>) on 2016-08-15.

Spontak, D. 2013. VACAPES MISSELEX Aerial Monitoring March 2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1017>) on 2016-08-15.

Spontak, D. 2014. Norfolk/VA Beach MINEX Vessel Surveys. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1072>) on 2016-08-15.

Spontak, D. 2015. Norfolk/VA Beach Inshore Vessel Surveys Nov 2012- Nov 2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1071>) on 2016-08-15.

Stevick, P. 2006. Allied Humpback Whale Catalogue, 1976 - 2003. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/73>) on 2016-08-15.

Stevick, P. 2013. YoNAH Encounter. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/274>) on 2016-08-15.

Swaim, Z. 2016. DUML vessel-based photo-id and biopsy surveys for proposed JAX USWTR site 2012-2015. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/906>) on 2016-08-15.

Swaim, Z. 2016. DUML vessel-based photo-id and biopsy surveys in Onslow Bay CHPT OPAREA 2011-2015. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/902>) on 2016-08-15.

Swaim, Z. 2016. DUML vessel-based photo-id and biopsy surveys in VACAPES OPAREA off Hatteras 2009, 2011-2015. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/907>) on 2016-08-15.

Taylor, J. 2015. Bottlenose dolphins off Outer Banks 2007-2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/837>) on 2016-08-15.

Thillet, M. 2011. Deep Panuke whale Acoustic 2003. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/651>) on 2016-08-15.

Thillet, M. 2011. Deep Panuke whale sightings 2003. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/650>) on 2016-08-15.

Tyson, R. 2014. Community structure and abundance of bottlenose dolphins *Tursiops truncatus* in coastal waters of the northeast Gulf of Mexico. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/571>) on 2016-08-15.

Urian, K. 2013. DUML New River surveys on the occurrence, distribution and density of marine mammals in Camp Lejeune 2010-2011. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/959>)

on 2016-08-15.

Urian, K. 2013. DUML surveys for the stock discrimination of bottlenose dolphins along the Outer Banks of North Carolina 2011-2012. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1010>) on 2016-08-15.

Urian, K. 2013. DUML vessel-based line transect surveys for proposed Onslow Bay USWTR site 2007-2010. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/433>) on 2016-08-15.

Urian, K. 2014. DUML coastal surveys on the occurrence, distribution and density of marine mammals in Camp Lejeune 2010-2013. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/957>) on 2016-08-15.

Van Parijs, S. 2013. NEFSC Marine Mammal Abundance Cruise 2004 Passive Acoustic Monitoring - Rainbow Click Detections. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/509>) on 2016-08-15.

Van Parijs, S. 2013. North Atlantic right whale up-calls in Stellwagen Bank National Marine Sanctuary 2006-2007. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/892>) on 2016-08-15.

Whitt, A. 2015. Marine mammal records of Cuba. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/1190>) on 2016-08-15.

Wolff, N. 2011. Aerial survey of upper trophic level predators on PLatts Bank, Gulf of Maine. Data downloaded from OBIS-SEAMAP (<http://seamap.env.duke.edu/dataset/103150267>) on 2016-08-15 and originated from iOBIS (<http://www.iobis.org>).